Content available at: https://www.ipinnovative.com/open-access-journals

# Journal of Dental Specialities

Journal homepage: https://www.jdsits.in/



## **Original Research Article**

# Comparative evaluation of coronally advanced flap using amniotic membrane and platelet rich fibrin in the treatment of Cairo's recession Type-I defects: A split mouth study

Deepmala Maurya<sup>1</sup>, Anju Rana<sup>2</sup>\*\*, Shweta Bali<sup>3</sup>, Meenu Taneja<sup>3</sup>, Priyanka Aggarwal<sup>3</sup>

#### **Abstract**

**Background:** The Coronally Advanced Flap (CAF) combined with Connective Tissue Graft (CTG) is the standard approach for managing gingival recession defects. However, it may cause increased patient discomfort due to the need for an additional surgical site. This study evaluates the efficacy of Amniotic Membrane (AM) and Platelet-Rich Fibrin (PRF) as alternatives for treating Cairo's Recession Type I defects.

Aim: To compare the clinical outcomes of CAF with AM (CAF+AM) and with PRF (CAF+PRF) in root coverage for Cairo's Recession Type I defects. **Materials and Methods:** Ten patients aged 20 to 40 years with bilateral Cairo's Recession Type I defects were enrolled. Sites were randomly assigned to receive either CAF+AM (test group) or CAF+PRF (control group). Clinical parameters including probing depth (PD), clinical attachment level (CAL), recession depth (RD), recession width (RW), gingival biotype (GB), width of keratinized tissue (WKT), and wound healing index (WHI)—were recorded at

on data distribution.

**Results:** Both groups showed statistically significant improvements in CAL, RD, RW, and WKT from baseline to 6 months (p < 0.05). However, the AM group demonstrated significantly greater root coverage (95.5% vs 77%; p < 0.01) and keratinized tissue gain. No significant differences were found between the groups for PD, GB, or WHI (p > 0.05). Healing was uneventful in both groups with good patient compliance and no adverse events.

baseline, and at 1, 3, and 6 months postoperatively. Statistical analysis was performed using t-tests, Wilcoxon signed-rank tests, and Friedman's test depending

**Conclusion:** Both AM and PRF in combination with CAF are effective in treating Cairo's Type I gingival recession defects. However, AM gave superior clinical outcomes in terms of root coverage and keratinized tissue width. These findings suggest AM may serve as a viable alternative to PRF. Further large scale and long term studies are warranted to validate these results and assess the stability of treatment outcomes.

Keywords: Amniotic membrane, Gingival recession, Guided tissue regeneration, Platelet rich fibrin

Received: 26-09-2024; Accepted: 07-08-2025; Available Online: 29-09-2025

This is an Open Access (OA) journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

# 1. Introduction

Miller characterizes gingival recession as the apical displacement of the gingival margin, resulting in the exposure of the root surface. Addressing this condition is particularly challenging for periodontists due to its sensitivity to technique and the subjective nature of the esthetic results perceived by patients. Gingival recession can stem from various causes, including bone dehiscence or fenestration defects, insufficient keratinized gingiva, and aberrant frenal

attachments. Contributing factors might include improper brushing techniques, dental plaque leading to periodontal inflammation, self-inflicted trauma, inappropriate orthodontic treatment, traumatic deep bites, and sub-gingival restorations. The multifaceted nature of these factors makes the treatment of gingival recession complex. Available treatment options include rotational pedicle flaps, advanced pedicle flaps, and both non-submerged and submerged grafts.

\*Corresponding author: Anju Rana Email: anjurana.sds@mrei.ac.in

<sup>&</sup>lt;sup>1</sup>Dept. of Periodontology, Maulana Azad Institute of Dental Sciences, New Delhi, Delhi, India

<sup>&</sup>lt;sup>2</sup>Dept. of Periodontology, Manav Rachna Dental College, School of Dental Sciences, MRIIRS, Faridabad, Haryana, India

<sup>&</sup>lt;sup>3</sup>Dept. of Periodontology, Santosh Dental College, Ghaziabad Uttar Pradesh, India

Careful planning is crucial to prevent esthetic dissatisfaction, increased clinical attachment loss, and root hypersensitivity. Cairo and Chambrone did a systematic review and concluded coronally Advanced Flap (CAF) along with Connective Tissue Graft (CTG) as the standard for gingival recession coverage.<sup>3-5</sup> Nevertheless, this method has limitations, comprising the prerequisite for an additional surgical site and amplified patient discomfort.<sup>3-5</sup> Alternatives to CTG, like Amniotic Membrane (AM) and Platelet-Rich Fibrin (PRF), have demonstrated comparable results. 6-8 PRF is particularly notable for its high levels of growth factors, platelets, and cytokines along with its excellent hemostatic properties, has been explored as a CTG viable substitute.<sup>6-8</sup> AM, a biological membrane from the chorion, contains an epithelial lining, a robust basement membrane, and avascular mesenchymal connective tissue, making it a suitable allograft. Literature suggests that AM, due to its fibronectins, proteoglycans, and collagen types, can achieve significant root coverage by closely resembling gingival tissue. 9-12 Nonetheless, there is limited evidence directly comparing AM with CTG, indicating a need for further research. While case reports and case series have highlighted the effectiveness of AM for treating gingival recession, a splitmouth study comparing these methods directly has yet to be conducted. 12-15 The present study aims to evaluate and compare the clinical efficacy of CAF with AM versus CAF with PRF for treating Cairo's Recession Type-I recession defects.16

## 2. Materials and Methods

This double-blinded randomized control trial was conducted in accordance with the Helsinki Declaration, with patient consent obtained and institutional ethics board approval secured. Armamentarium used is shown in **Figure 1**. The study included 10 participants, aged 20 to 40 years, who met the inclusion criteria of being free from non-carious cervical lesions, not having undergone periodontal surgery in the past 12 months, and having no local predisposing factors, smoking history, lactation, or systemic diseases that could affect treatment outcomes.<sup>17</sup>

Baseline measurements were collected (**Figure 2**) after basic oral prophylaxis, including sulcus probing depth (PD), gingival biotype (GB), clinical attachment level (CAL), recession depth (RD), width of recession (RW), and keratinized gingiva width (WKT). <sup>17,18</sup> A split-mouth design was employed, with bilateral Cairo's Type-I recession defects treated using CAF+AM (test group) (**Figure 3**) and CAF+PRF (control group), (**Figure 4**) assigned through computer-assisted randomization. Patients were blinded to procedure being performed on each site.

These parameters were assessed at baseline, 1 month and every three months upto 6 months postoperatively. Additionally, Index of Wound Healing (WHI) was recorded post-surgery, with scores assigned as follows: Score 1 for uneventful healing, Score 2 for slight complications, and

Score 3 for poor healing. Measurements were standardized using a UNC-15 periodontal probe, a stent, and a vernier caliper.


The CAF procedure involved horizontal and vertical incisions, followed by root planing and EDTA treatment. PRF was prepared by centrifuging patient blood, while AM was cut to size and applied to the recession site. The gingival flaps were coronally advanced, secured with interrupted sutures, and covered with a periodontal pack. Postoperative care included systemic analgesics, 0.2% chlorhexidine gluconate, and the Charters brushing technique. Statistical analyses were performed using the independent t-test, Wilcoxon signed-rank test, and Friedman's test, depending on data distribution.

#### 3. Results


In this split-mouth randomized control trial, 10 subjects with 20 bilateral cairo's type I recession defects sites were treated with either AM combined with CAF or PRF combined with CAF. Single calibrated examiner recorded the records of all participants included in the study. Various clinical parameters recorded by the calibrated examiner were PD, RD, WKT, GB, CAL, RW and WHI. All these parameters were re-examined at 1, 3, and 6 months (Figure 5). Table 1 describes the gender distribution in the study and Table 2 depict the site involved. There were eight males (80%) and two females (20%) among the ten patients involved. **Table 3** explains the parameters recorded. As indicated in Table 3, there were no statistically significant changes in probing depth (PD), gingival biotype (GB), and wound healing index (WHI) from baseline to 6 months. Though, significant alterations were detected in clinical attachment level (CAL) (p=0.045, 0.034, 0.014 for 1, 3, and 6 months, respectively), recession depth (RD) (p=0.031, 0.016, 0.001), recession width (RW) (p=0.035, 0.001, 0.007), and width of keratinized tissue (WKT) (p=0.042, 0.001) at the 3 and 6-month marks. Table 4 describes an analysis of the mean root coverage percentages amongst the control and test groups during the follow-up period. Significant differences were observed supporting the test group (p=0.001, 0.003, and 0.001) at 1, 3, and 6 months, respectively.



Figure 1: A: Armamentarium used; B: Customized acrylic stent




**Figure 2:** Clinical measurements recorded before surgical procedure; **A:** Recession height (RH); **B:** Recession width (RW); **C:** Probing depth (PD); **D:** Width of keratinized tissue (WKT); **E:** Gingival biotype (GB)



**Figure 3:** Coronally advanced flap with amniotic membrane (test site); **A:** Incision design; **B:** Flap elevation and coronal advancement; **C:** Air dried amniotic membrane; **D:** Placement of membrane; **E:** Suturing



**Figure 4:** Coronally advanced flap with amniotic membrane (Control site); **A:** Incision design; **B:** Flap elevation and coronal advancement; **C:** Coronal advancement of flap with PRF placement; **D:** Suturing



**Figure 5:** Follow up for test and control group at baseline, 3 month and 6 months; **A:** At baseline (Test group); **B:** At baseline (Control group); **C:** Post-operative 3 month (Test group); **D:** Post-operative 3 month (control group); **E:** Post-operative 6 month (Test group); **F:** Post-operative 6 month (control group)

**Table 1:** Gender distribution of patients included into the study.

	Number	Percentage
Male	8	80.0%
Female	2	20.0%
Total	10	100%

**Table 2:** Distribution of sites for the surgical technique performed.

Tooth Number	Groups			
	Test group	Control group		
13	1	1		
	10.0%	10.0%		
23	1	1		
	10.0%	10.0%		
33	6	2		
	60.0%	20.0%		
43	2	6		
	20.0%	60.0%		
Total	10	10		
	100.0%	100.0%		

**Table 3:** Showing the intergroup comparisons of various parameters evaluated during the 6- month follow-up (p Value < 0.05 was considered as statistically significant).

		Test group		Control group		P value
	Mean	SD	Mean	SD		
PD	•			•		
Baseline	1.50	0.53	1.40	0.48	0.20	0.388
At 1 month	1.50	0.53	1.40	0.48	0.20	0.388
At 3 months	1.50	0.53	1.40	0.48	0.20	0.388
At 6 months	1.50	0.53	1.40	0.48	0.20	0.388
CAL	•	1		•	<u>'</u>	
Baseline	4.50	0.97	4.40	0.70	0.10	0.795
At 1 month	2.98	0.71	3.42	0.72	-0.44	0.045*
At 3 months	2.21	0.58	2.73	0.72	-0.52	0.034*
At 6 months	1.44	0.41	2.05	0.57	-0.61	0.014*
RD	•			•		
Baseline	2.90	0.88	2.90	0.57	0.00	1.000
At 1 month	1.48	0.56	2.01	0.57	-0.53	0.031*
At 3 months	0.71	0.43	1.27	0.51	-0.56	0.016*
At 6 months	0.16	0.24	0.68	0.37	-0.52	0.001*
RW	<u>.</u>			•		
Baseline	3.13	0.46	3.12	0.38	0.01	0.958
At 1 month	1.73	0.45	2.03	0.41	-0.30	0.035*
At 3 months	0.52	0.39	1.18	0.40	-0.66	0.001*
At 6 months	0.21	0.25	0.59	0.30	-0.38	0.007*
WKT	<u>.</u>			•		
Baseline	0.98	0.43	1.00	0.45	-0.02	0.920
At 1 month	1.98	0.53	1.86	0.37	0.12	0.562
At 3 months	3.03	0.32	2.72	0.34	0.31	0.042*
At 6 months	3.76	0.33	3.15	0.35	0.61	0.001*
GB	<u>.</u>			•		
Baseline	1.22	0.32	1.29	0.20	-0.07	0.574
At 1 month	1.30	0.22	1.29	0.23	0.01	0.922
At 3 months	1.25	0.20	1.27	0.22	-0.02	0.833
At 6 months	1.28	0.15	1.31	0.17	-0.03	0.688
WHI	·			•		
Baseline	1.20	0.42	1.30	0.48	-0.10	0.628
At 1 month	1.00	0.00	1.00	0.00	0.00	1.000
At 3 months	1.00	0.00	1.00	0.00	0.00	1.000
At 6 months	1.00	0.00	1.00	0.00	0.00	1.000

<sup>\*</sup>Statistically significant

Std. Percentage root Groups Mean Mean t-test pcoverage **Deviation** difference value value 1 month Test group 49.83 10.44 3.80 3.799 0.001\* Control group 31.33 11.33 77.17 3 months Test group 12.82 3.48 3.484 0.003\* 57.00 Control group 13.07 95.50 4.29 4.294 0.001\* Test group 6.66 6 months Control group 77.33 11.61

**Table 4:** Shows the intergroup comparison of percentage of root coverage evaluated during the 6- month follow-up (p Value < 0.05 was considered as statistically significant and p value  $\le 0.001$  was considered as statistically highly significant).

#### 4. Discussion

This split-mouth randomized control, double-blind clinical study was conducted to assess the clinical efficacy of AM combined with CAF technique for treating Cairo's Type-I defects. At baseline, there were no statistically significant differences between the groups evaluated in any of the measured clinical parameters. Bleeding on probing and plaque presence were recorded at all follow-ups to monitor oral hygiene and inflammation, and all patients maintained good oral hygiene, with no signs of inflammation or bleeding on probing.

Throughout the study, no significant changes in probing depth (PD) (p>0.05) were observed from baseline to 6 months in either group, consistent with findings from Agarwal et al.<sup>9</sup> and Gautam.<sup>19</sup> However, Wallace et al.<sup>20</sup> reported a 0.8 mm reduction in PD over the same period when comparing amniotic membrane to acellular dermal matrix.

Both the AM and PRF groups showed significant improvements in clinical attachment level (CAL) from baseline to 6 months, indicating effective root coverage by both techniques. Nonetheless, a significant difference was noted in CAL gain between the groups. AM's composition, resembling the basement membrane of oral mucosa and containing laminin 5, may contribute to improved gingival cell adhesion and CAL enhancement, as highlighted by Gurinsky,<sup>21</sup> Meller et al.,<sup>22</sup> and Riau et al.<sup>23</sup>

Our study found a greater percentage of root coverage in the AM group (95.5%) compared to the PRF group (77%), contrasting with Agarwal et al., 10 who reported greater root coverage with PRF. Studies by Gurinsky, 21 Agarwal et al., 9 and Gautam 19 showed varying levels of root coverage with AM, while Aroca et al., 24 Jankovic et al., 8 and Agarwal et al. 9 reported diverse results with PRF. Despite AM's superior root coverage in this study, neither method achieved 100% root coverage.

Significant improvement in recession width (RW) was observed at 3 and 6 months in both groups, similar to findings by Atilla et al.,<sup>25</sup> Aroca et al.,<sup>24</sup> and Uraz et al.<sup>26</sup> for PRF-treated sites. Agarwal et al.<sup>9</sup> noted a greater reduction in RW for PRF at 3 months, but similar values at 6 months, with AM

showing increased RW. The preservation of the mucogingival junction and granulation tissue from the periodontal ligament might enhance keratinized gingiva. Agarwal et al. 9 and Gautam 19 reported significant increases in attached gingiva (AG) with PRF and AM, though PRF showed more width increase. Other studies by Jankovic et al. 8 and Uraz et al. 26 noted greater increases in keratinized tissue width with PRF compared to SCTG, while Wallace et al. 20 reported a 0.2 mm increase with AM at 4 months.

Gingival biotype improvement was statistically significant in both groups from baseline to 6 months, with no significant difference between them (p>0.05), suggesting both materials effectively enhance gingival thickness. This finding is consistent with Shetty et al.,<sup>27</sup> though Agarwal et al.<sup>9</sup> found more substantial thickness increases with PRF compared to AM.

The Wound Healing Index (WHI) did not show significant differences between the groups (p>0.05) at 6 months, indicating uneventful healing for both treatments. This aligns with the properties of both materials used.

Agarwal et al.9 also assessed patient satisfaction, finding higher comfort scores for AM and better esthetic scores for PRF. Based on our results, AM appears superior to PRF for treating gingival recession, though neither method achieved complete root coverage. Variations in clinical parameter can measurement studies methods among comparisons. The study's limitations include a small sample size, short duration, and lack of histological evaluation due to ethical constraints. Also patient centered outcome like pain assessment was not taken into consideration. Larger, longterm, multicenter randomized controlled trials with patient centered outcomes are needed to further evaluate AM's efficacy compared to PRF.

# 5. Conclusion

Based on the findings, it can be concluded that both AM and PRF, when used with CAF technique, are effective treatments for Cairo's recession type-I defects. Both materials effectively reduced recession width and depth, and augmented width of keratinized tissue. They also demonstrated positive outcomes in improving gingival

<sup>\*</sup>Statistically significant

thickness and facilitated good and rapid healing. However, AM provided greater root coverage and better keratinized tissue width compared to PRF. Therefore, AM may be considered a superior option over PRF for treating recession defects. Nonetheless, additional research is needed to evaluate the long term stability of the tissue thickness and root coverage achieved. Future studies with larger sample sizes and longer follow-up periods are necessary to further confirm AM's superior efficacy compared to PRF.

# 6. Source of Funding

None.

### 7. Conflict of Interest

None.

#### References

- Tarnow D, Hochman M, Chu S, Fletcher P. A New Definition of Attached Gingiva Around Teeth and Implants in Healthy and Diseased Sites. Int J Periodontics Restor Dent. 2021;41(1):43–9.
- Periodontology AAO. Glossary of periodontal terms. 4th edn. American Academy of Periodontology. Chicago, Illinois. 2001
- Cairo F, Pagliaro U, Nieri M, Treatment of gingival recession with coronally advanced flap procedures: a systematic review. *J Clin Periodontol*. 2008;35(Supplement 8):136–62.
- Chambrone L, Tatakis DN. Periodontal soft tissue root coverage procedures: a systematic review from the AAP Regeneration Workshop. J Periodontol. 2015;86(2 Suppl):S8–51.
- Cairo F, Nieri M, Pagliaro U. Efficacy of periodontal plastic surgery procedures in the treatment of localized facial gingival recessions. A systematic review. J Clin Periodontol. 2014;41(Suppl 15):S44– 62.
- Padma R, Shilpa A, Kumar PA, Nagasri M, Kumar C, Sreedhar A.
   A split mouth randomized controlled study to evaluate the adjunctive effect of platelet-rich fibrin to coronally advanced flap in Miller's class-I and II recession defects. *J Indian Soc Periodontol*. 2013;17(5):631.
- Dohan DM, Choukroun J, Diss A, Dohan SL, Dohan AJJ, Mouhyi J, Gogly B, et al., Platelet-rich fibrin (PRF): a second-generation platelet concentrate. Part II: platelet-related biologic features. *Oral Surg Oral Med Oral Pathol Oral Radiol Endodontol*. 2006;101(3):e45–e50.
- Jankovic S, Aleksic ZM, Klokkevold PR, Lekovic V, Dimitrijevic B, Kenney EB, et al. Use of platelet-rich fibrin membrane following treatment of gingival recession: a randomized clinical trial. *Int J Period Restor Dent.* 2012;32(2):165.
- Agarwal SK, Jhingran R, Bains VK, Srivastava R, Madan R, Rizvi I. Patient-centered evaluation of microsurgical management of gingival recession using coronally advanced flap with platelet-rich fibrin or amnion membrane: A comparative analysis. *Eur J Dent*. 2016;10(01):121–33.
- Bolla V, Reddy PK, Kalakonda B, Koppolu P, Manaswini E. Coronally advanced flap with amniotic membrane in the treatment of gingival recession: Three case reports. *Int J Appl Basic Med Res*. 2019;9(2):111–4.
- Ghahroudi A, Khorsand A, Rokn AR, Sabounchi SS, Shayesteh YS, Soolari A. Comparison of amnion allograft with connective tissue graft for root coverage procedures: a double-blind,

- randomized, controlled clinical trial. *J Int Acad Periodontol*. 2013;15(4):101–12.
- Shah R, Sowmya N, Mehta D, Amnion membrane for coverage of gingival recession: A novel application. *Contemp Clin Dent*. 2014;5(3):293–5.
- Singh H, Singh H. Bioactive amnion as a guided tissue regeneration (GTR) membrane for treatment of isolated gingival recession. A case report. *Indian J Dent.* 2013;4(2):110–13.
- Mahajan R, Gill AS, Khinda PK, Shewale A, Saravanan SP. Guided tissue regeneration based treatment of root coverage using placental membrane allograft: A case report. *IJSS Case Rep Rev.* 2015;2:1–5.
- Sharma A, Yadav K. Amniotic membrane-A Novel material for the root coverage: A case series. *J Indian Soc Periodontol*. 2015;19(4):444–8.
- Cairo F, Nieri M, Cincinelli S, Mervelt J, Pagliaro U. The interproximal clinical attachment level to classify gingival recessions and predict root coverage outcomes: an explorative and reliability study. J Clin Period. 2011;38(7):661–6.
- Kasaj A. Definition of Gingival Recession and Anatomical Considerations. In: Kasaj, A. (eds) Gingival Recession Management. Springer, Cham. 2018. https://doi.org/10.1007/978-3-319-70719-8
- De Sanctis M, Zucchelli G. Coronally advanced flap: A modified surgical approach for isolated recession-type defects: Three-year results. J Clin Periodontol. 2007;34(3):262–8.
- Gautam A. Comparative evaluation of coronally advanced flap with platelet-rich fibrin vs amniotic membrane for the treatment of gingival recession. J Health Sci Res. 2017;8(1):31–7.
- Wallace SC. Root coverage grafting comparing placental derived membrane to acellular dermis matrix: A case series. *Dentistry*. 2012;2(137):1-4.
- Gurinsky B. A novel dehydrated amnion allograft for use in the treatment of gingival recession: An observational case series. J Implant Adv Clin Dent. 2009;1(1):65–73.
- Meller D, Pires RT, Mack RJ, Figueiredo F, Heiligenhaus A, Park WC, et al., Amniotic membrane transplantation for acute chemical or thermal burns. *Ophthalmology*. 2000;107(5):980–9.
- Riau AK, Beuerman RW, Lim LS, Mehta JS. Preservation, sterilization and de-epithelialization of human amniotic membrane for use in ocular surface reconstruction. *Biomaterials*. 2010;31(2):216–25.
- Aroca S, Beuerman RW, Lim LS, Mehta JS. Clinical evaluation of a modified coronally advanced flap alone or in combination with a platelet-rich fibrin membrane for the treatment of adjacent multiple gingival recessions: A 6-month study. *J Periodontol*. 2009;80(2):244–52.
- Eren G, Atilla G. Platelet-rich fibrin in the treatment of localized gingival recessions: a split- mouth randomized clinical trial. *Clin Oral Investig.* 2014;18(8):1941–8.
- Uraz A, Bains VK, Jhingran R, Srivastava R, Mada R n. Comparative evaluation of platelet-rich fibrin membrane and connective tissue graft in the treatment of multiple adjacent recession defects: a clinical study. J Dent Sci. 2015;10(1)::36–45.
- Shetty SS, Chatterjee A, Bose S. Bilateral multiple recession coverage with platelet-rich fibrin in comparison with amniotic membrane. *J Indian Soc Periodontol*. 2014;18(1):102.

Cite this article: Maurya D, Rana A, Bali S, Taneja M, Aggarwal P. Comparative evaluation of coronally advanced flap using amniotic membrane and platelet rich fibrin in the treatment of Cairo's recession Type-I defects: A split mouth study. *J Dent Spec*. 2025;13(2):220-225.